

Production of C-glucosides of flavonoids and related compounds by *Escherichia coli* expressing buckwheat C-glucosyltransferase

Takamitsu Ito, Shunsuke Fujimoto, Makoto Shimosaka, Goro Taguchi*

Faculty of Textile Science and Technology, Shinshu University, Ueda, Nagano 386-8567, Japan

*E-mail: gtagtag@shinshu-u.ac.jp Tel: +81-268-21-5342 Fax: +81-268-21-5331

Received September 18, 2014; accepted October 16, 2014 (Edited by A. Sugiyama)

Abstract C-Glucosides are glucose-containing glycosides that have carbon–carbon bonds between the anomeric carbon of the sugar moieties and aglycon, rendering the molecules remarkably stable against hydrolysis by enzymes or acids. In this work, we showed the production of C-glucosides of flavonoids and related compounds (i.e., 2-hydroxyflavanone, dihydrochalcone, and trihydroxyacetophenone) by *Escherichia coli* expressing buckwheat C-glucosyltransferase. The substrates in their respective cultures were taken up by the cells and C-glucosylated, and the products were released into the culture media. The bioconversion process was completed in 1–2h, but products were already observed immediately after addition of the substrates (200 μ M). The conversion rates of these substrates reached 80–95%. Without addition of glucose to the conversion media, almost no C-glucosides were produced. Although the amounts of the substrates fed to their respective cultures were limited by their solubility in water, repeated addition of the substrate to the culture at regular time intervals effectively increased the total amount of product obtained.

Key words: Flavonoid C-glucoside, bioconversion in *E. coli*, C-glucoside production, C-glucosyltransferase, biocatalyst.

Introduction

Glycosylation is one of the predominant processes used by plants to modify their secondary metabolites. The process usually occurs at O-, N-, S-, or C-atoms of the metabolites, with the most common plant glycosides being O-glucosides (Jones and Vogt 2001). With this modification, the compounds possess increased water solubility and chemical stability, which regulate their biological activity and accumulation (Bowles et al. 2005; Gachon et al. 2005). Thus, glycosylation is important for the management of secondary metabolites in plants. With the exception of some anthocyanin glycosylations (Matsuba et al. 2010), glycosylations are usually catalyzed by uridine 5'-diphosphate (UDP)-sugar: glycosyltransferases (UGTs) (Bowles et al. 2005; Gachon et al. 2005).

C-Glucosides are characterized by the carbon–carbon bond that forms between the anomeric carbon of sugar moieties and the carbon atom of aglycon (Franz and Grün 1983). Because of this bond, C-glycosides are extremely stable against enzyme or acid hydrolysis

compared with other glycosides that can be hydrolyzed. Many kinds of plant C-glycosides have been reported (Franz and Grün 1983), and the most common are the flavonoid C-glycosides that have been found in mosses, liverworts, ferns, gymnosperms, monocots, and dicots (Jay 1994; Talhi and Silva 2012). These C-glycosides are known to show various biological activities in plants, such as antioxidant, antifungal, insecticide, and allelochemical activities (Byrne et al. 1996; Hooper et al. 2010; Jay 1994; McNally et al. 2003). In animals, flavonoid C-glycosides have shown antioxidant, anti-inflammatory, antihyperglycemic, anti-obesity, and anti-ulcer activities (reviewed by Talhi and Silva 2012). Recently, many other medicinal effects were also reported, such as the abilities to potentiate pentobarbital-induced sleep (Wang et al. 2008) and to confer beneficial effects on memory loss (de Oliveira et al. 2014; Jung et al. 2014). Thus, flavonoid C-glycosides have received much attention for their human health and pharmaceutical applications.

Flavonoid C-glycosides are obtained by purification from natural resources. However, this is difficult and

Abbreviations: CGT, C-glucosyltransferase; DMSO, dimethyl sulfoxide; HMBC, heteronuclear multiple bond correlation; HPLC, high-performance liquid chromatography; LC-MS, liquid chromatography-mass spectrometry; NMR, nuclear magnetic resonance; PhTHAP, 2-phenyl-2',4',6'-trihydroxyacetophenone; THAP, 2',4',6'-trihydroxyacetophenone; UDP-glucose, uridine 5'-diphosphoglucose; UGT, UDP-sugar: glycosyltransferase. This article can be found at <http://www.jspcmb.jp/>

Published online January 14, 2015




Figure 1. Substrates used in this study.

expensive in many cases, because these compounds are present at low concentrations and the process is complicated by the removal of other constituents. Chemical syntheses of flavonoid C-glucosides have been reported (reviewed by Talhi and Silva 2012), but the processes required many steps and resulted in relatively low total yields.

Brazier-Hicks et al. (2009) isolated a C-glucosyltransferase (CGT) gene from rice (OsCGT) and characterized it as belonging to the UGTs. The substrates of CGT were 2-hydroxyflavanones, and the products were then dehydrated to become C-glucosylflavones. Using this OsCGT enzyme, two groups have reported the production of flavone- and/or dihydrochalcone-C-glucosides via biotransformation in the yeast expression system (Brazier-Hicks and Edwards 2013) and enzymatic conversion coupled with sucrose synthase (Bungaruang et al. 2013).

We recently characterized CGTs from buckwheat (FeCGTa and FeCGTb; Nagatomo et al. 2014). These CGTs showed significant activity against flavonoid substrates possessing a basic skeleton of 2',4',6'-trihydroxyacetophenone (THAP), such as 2-hydroxyflavanones and dihydrochalcone. They also reacted significantly with non-flavonoid substrates possessing THAP-like skeleton, such as THAP and 2-phenyl-2',4',6'-trihydroxyacetophenone (PhTHAP). Compare to these flavonoid substrates, THAP has a structure lacking B-ring of flavonoids, and PhTHAP has a truncated methylene chain between A- and B-ring (Figure 1). OsCGT showed only faint activities against THAP, suggesting the broader substrate specificity of the FeCGTs. Therefore, FeCGTs were expected to be good tools for the biological production of C-glucosides.

In this study, we applied an *Escherichia coli* expression system for the production of C-glucosides, as an easy and useful biocatalyst. Lim and colleagues (Lim 2005; Lim et al. 2004) proposed *E. coli* expressing the plant UGT genes as novel biocatalysts for the bioconversion of aglycons to their glycosides, and other authors (Kim et al. 2012, 2013) have achieved high production yields of glycosides

with these expression systems.

Materials and methods

Reagents

The reagents used were as follows: phloretin and THAP (Tokyo Chemical Industries, Tokyo, Japan), PhTHAP (Sigma-Aldrich, St. Louis, MO, USA), apigenin (Indofine Chemicals, Hillsborough, NJ, USA), chrysin and DMSO-*d*₆ (Across Organics, Geel, Belgium), vitexin and isovitexin (Extrasynthèse, Genay, France). 2-Hydroxyypinocembrin and 2-hydroxynaringenin were synthesized from chrysin and apigenin, respectively, as previously described (Nagatomo et al. 2014). All other chemicals were obtained from Wako Pure Chemical Industries, Ltd. (Osaka, Japan), Nacalai Tesque (Kyoto, Japan), and Kanto Chemical Co., Inc. (Tokyo, Japan), unless otherwise specified.

Bioconversion of phenolic substrates by *E. coli* expressing recombinant FeCGTa

FeCGTa (UGT708C1; GenBank Accession No. AB909375) cDNA subcloned into pET28a(+) (Merck, Darmstadt, Germany) was transformed into *E. coli* Rosetta™ 2(DE3) (Merck), as described previously (Nagatomo et al. 2014). An aliquot (1 ml) of overnight culture was added to 200 ml of LB medium (Nacalai Tesque) containing kanamycin (50 mg l⁻¹) and chloramphenicol (34 mg l⁻¹), and this culture was incubated at 37°C with shaking at 150–200 rpm until the OD₆₀₀ reached 0.6. Isopropyl-β-D-1-thiogalactopyranoside (final concentration, 0.4 mM) was added and the culture was further incubated at 22°C for 24 h with shaking at 150–200 rpm. The bioconversion was performed according to the method described by Kim et al. (2012) with some modifications. The bacterial cells expressing FeCGTs were precipitated at 4000×*g* for 10 min, and the pellet was suspended in M9 basal medium (Green and Sambrook 2012) containing 2% glucose, adjusting the cell density to an OD₆₀₀ value of 3.0–6.0. This suspension was used for the conversion of the compounds. Substrates were dissolved in DMSO at 100–200 mM and added to the suspension at a final concentration of 200 μM. The culture (3 ml for small scale, 50–200 ml for production of C-glucosides) was incubated

at 30°C with shaking at 150–200 rpm, and an aliquot of the medium was collected at appropriate intervals for product determination. To confirm an effect of sugar concentration on the conversion efficiency, glucose (0.4–2%) or maltose (2%) was added into M9 basal medium, and phloretin was used as a substrate. For production from the substrates phloretin and PhTHAP, the substrate was added at six different times to 200 ml of culture, at 1.5–2.5 h intervals, for a total of 12 h. The culture medium containing the compounds was adsorbed to a Sep-Pak Plus C18 column (Waters, Milford, MA, USA) to remove residual glucose, and was then eluted by methanol. The eluates were diluted with water and adsorbed to an ODS column (Wako-gel 50C18, 15 mm i.d. × 120 mm; Wako Pure Chemical Industries, Ltd.), equilibrated with 20% methanol, for purification. The compounds were eluted with a linear gradient of 20–60% methanol (300 ml), and the fraction containing the product was concentrated and crystallized. NMR spectra were recorded on a Bruker Avance400 spectrometer (Bruker Biospin, Yokohama, Japan), and compared with reported NMR spectra of related compounds (Ogawa et al. 2001; Yepremyan et al. 2010). Phloretin C-glucoside (nothofagin): ^1H NMR (400 MHz, $\text{DMSO-}d_6$): δ 13.77 (1H, s, Ar-OH), 11.03 (1H, s, Ar-OH), 10.05 (1H, s, Ar-OH), 9.12 (1H, s, Ar-OH), 7.02 (2H, d, $J=8.4$ Hz, H-2 and H-6), 6.66 (2H, d, $J=8.4$ Hz, H-3 and H-5), 5.94 (1H, s, H-5'), 4.82 (1H, m, glucose-OH), 4.79 (1H, m, glucose-OH), 4.52 (1H, d, $J=9.8$ Hz, H-1''), 4.48 (2H, m, glucose-OH), 3.88 (1H, m, H-2''), 3.66 (1H, m, H-6''), 3.42 (1H, m, H-6''), 3.22 (2H, t, $J=8.2$ Hz, H- α), 3.14 (3H, m, H-3''–H-5''), 2.78 (2H,

$J=8.0$ Hz, H- β); ^{13}C NMR (100 MHz, $\text{DMSO-}d_6$): δ 204.8 (C=O), 165.1 (C-2'), 164.0 (C-4'), 162.0 (C-6'), 155.7 (C-4), 132.0 (C-1), 129.5 (C-2 and C-6), 115.4 (C-3 and C-5), 104.4 (C-3'), 104.0 (C-1'), 95.0 (C-5'), 81.7 (C-5''), 79.3 (C-3''), 73.9 (C-1''), 71.0 (C-2''), 70.8 (C-4''), 61.6 (C-6''), 46.0 (C- α), 29.9 (C- β). 2-Phenyl-3'-C-glucosyl-2',4',6'-trihydroxyacetophenone (PhTHAP-C-glucoside): ^1H NMR (400 MHz, $\text{DMSO-}d_6$): δ 13.76 (1H, s, Ar-OH), 11.08 (1H, s, Ar-OH), 10.12 (1H, s, Ar-OH), 7.31–7.19 (5H, m, H-2–H-6), 5.96 (1H, s, H-5'), 4.81 (2H, m, glucose-OH), 4.51 (1H, d, $J=9.8$ Hz, H-1''), 4.47 (2H, m, glucose-OH), 4.36 (2H, s, H- α), 3.89 (1H, m, H-2''), 3.65 (1H, d, $J=11.5$ Hz, H-6''), 3.42 (1H, d, $J=10.9$ Hz, H-6''), 3.16–3.12 (3H, m, H-3''–H-5''); ^{13}C NMR (100 MHz, $\text{DMSO-}d_6$): δ 202.9 (C=O), 165.4 (C-2'), 164.4 (C-6'), 161.9 (C-4'), 136.4 (C-1), 130.0 (C-2 and C-6), 128.5 (C-3 and C-5), 126.6 (C-4), 104.5 (C-3'), 103.9 (C-1'), 95.0 (C-5'), 81.7 (C-5''), 79.3 (C-3''), 73.9 (C-1''), 71.0 (C-2''), 70.9 (C-4''), 61.7 (C-6''), 49.4 (C- α).

LC-MS analysis

LC-MS analysis was performed using a Waters UPLC ACQUITY SQD system (Waters) with an electron-spray ionization probe, as described previously (Nagatomo et al. 2014).

Results and discussion

C-Glycosides have received much attention for their human health and pharmaceutical applications, owing




Figure 2. Production of C-glycosides of flavonoids and related compounds by *Escherichia coli* expressing buckwheat C-glucosyltransferase (FeCGTa). A. Changes in the culture media after the addition of phloretin (final, 200 μM). Filled circles and open circles represent the concentrations of phloretin and nothofagin, respectively. B. HPLC profiles of the culture media at 0 and 1.5 h after the addition of phloretin. Peak identifications: 1, phloretin; 2, nothofagin; 3, uncharacterized metabolite of nothofagin. C. Mass spectrum of peak 2 (2.7 min) from B. D. HPLC profiles of culture media at 0 and 1.5 h after the addition of substrates. a: 2-Hydroxynaringenin; b: 2-hydroxypinocembrin; c: 2-phenyl-2',4',6'-trihydroxyacetophenone (PhTHAP); and d: 2',4',6'-trihydroxyacetophenone (THAP). Black and gray arrowheads indicate the peaks of administered substrates and their C-glycosides, respectively. The small peaks in "a" are the dehydrated products of C-glycoside (vitexin and isovitexin).

to their outstanding stability against enzyme and acid hydrolyses. We attempted to produce the C-glucosides of flavonoids and related compounds, using an *E. coli* strain expressing FeCGTa (Ec-FeCGT). The FeCGTa enzyme catalyzes the C-glucosylation of the open-circular form of 2-hydroxyflavanone and compounds possessing a THAP-like structure, shown in Figure 1 (Nagatomo et al. 2014). The substrate dihydrochalcone phloretin ($200\ \mu\text{M}$) was added to the Ec-FeCGT culture, and the medium contents were monitored at appropriate intervals. HPLC analysis revealed that most of the phloretin (Figure 2B Peak 1) had disappeared after its addition to the culture, and a more hydrophilic compound appeared in the medium (Peak 2). LC-MS analysis identified the compound as the C-glucoside of phloretin (nothofagin) (Figure 2C), showing both the $[\text{M}-\text{H}]^-$ ion at m/z 435 and the fragment ion $[\text{M}-\text{H}-120]^-$ at m/z 315 that is typical of C-glucosides (Jay 1994). Other substrates such as 2-hydroxyflavanones and THAP-like compounds (shown in Figure 1) were also added to the Ec-FeCGT system, and were converted to their C-glucosides and released into the media (Figure 2D). The conversions were fast, such that we detected the products even in the 0 h samples, because it took 1–2 min for the cells and the medium to separate after addition of the substrate. The conversion was mostly completed in 1–2 h for all the compounds tested at this condition, and the conversion rate of these compounds reached 80–95% (Figure 2). In the conversion of phloretin, when the incubation was continued even after the substrate had been expended, a small amount of nothofagin was sometimes modified further to produce an unknown compound (Figure 2B peak 3). The compound exhibited the $[\text{M}-\text{H}]^-$ ion at m/z 507, which was bigger than that of nothofagin at m/z 72. This modification was also observed in the conversions of THAP and PhTHAP. We have no idea of the structure of this modification at present. The effect of glucose on the conversion efficiency was tested using phloretin as a substrate (Figure 3). At the glucose concentration of 0.4–2%, most of the phloretin had converted to nothofagin within 1.5 h. In contrast, in the absence of glucose, almost no C-glucosides were found in the culture media. When maltose (a dimer of glucose that can be used as a sugar source by *E. coli*) was added at the concentration of 2%, the conversion rate was significantly low. These results showed that the addition of glucose was important for the bioconversion, and 0.4% of glucose was enough for the conversion, with a substrate concentration at $200\ \mu\text{M}$. The glucose could work as a source of the glucose moiety of UDP-glucose, and as an energy source for the production of UDP-glucose.

We then determined whether repeated substrate additions to the culture medium would result in the production of large amounts of the C-glucosides of each substrate tested. The result of repeated phloretin feeding




Figure 3. Effect of glucose concentration on C-glucoside production by *Escherichia coli* expressing buckwheat C-glucosyltransferase (FeCGTa). Glc, glucose; Mal, maltose.




Figure 4. Production of C-glucoside by *Escherichia coli* expressing buckwheat C-glucosyltransferase (FeCGTa), with repeated addition of the substrate. Phloretin ($200\ \mu\text{M}$) was added to the culture of *E. coli* expressing FeCGTa in 1.5–2 h intervals, as indicated in the figure, and analyzed by HPLC as described in the “Materials and methods” section. Black and gray arrowheads indicate the peaks of added phloretin and produced nothofagin, respectively.

(total 102 mg) is shown in Figure 4. At every 1.5–2.5 h, the fed phloretin (retention time at 3.35 min) had mostly disappeared from the media, and the peak of nothofagin (retention time at 2.6 min) increased instead. The total conversion efficiency was more than 90%. Repeated feeding until 9 h was effective for the bioconversion of the substrate. After purification by column chromatography, 110 mg of nothofagin was obtained (67% yield from phloretin). In the case of PhTHAP, repeated feeding resulted in an approximately 70% conversion to its C-glucosides. Both of the products were purified and analyzed by NMR in $\text{DMSO}-d_6$. The correlation between H-1 of glucose (δ 4.52 and 4.51) and C-3' of the compounds (δ 104.0 and 103.9) was observed by HMBC

analysis, which proved that these products were indeed C-glucosides.

The production of C-glucosides by means of bioconversion using OsCGT has been reported by two groups. Brazier-Hicks and Edwards (2013) reported an 80% yield of nothofagin from 500 μ M phloretin in 24 h, in a yeast expressing OsCGT. They also reported a high yield of C-glucosyl-2-hydroxynaringenin from naringenin that was converted by a yeast polyprotein system co-expressing a flavanone 2-hydroxylase from rice and an OsCGT. Compared with the yeast system, the conversion rate by our Ec-FeCGT system was significantly faster, converting more than 90% of phloretin (200 μ M) in 1–2 h (Figure 2A). The fed substrates are supposed to enter into *E. coli* cells by simple diffusion, so that an incorporation rate would be affected by a difference of their concentrations between inside and outside of the cells. Therefore, a high intracellular glucosyltransferase activity is favorable to keep the incorporation rate constant, because the incorporated substrates are efficiently glucosylated and successively excreted out of the cells. In general, a yield of recombinant proteins in *E. coli* is higher than that in budding yeast, which would have resulted in a fast conversion rate of the substrate in Ec-FeCGT system. It should be noted that a suitable concentration of the fed substrates is important for an efficient conversion. When phloretin was added to the medium at an initial concentration of 2 mM, a precipitation of the substrate occurred, and a conversion to nothofagin was scarcely observed (data not shown). Our Ec-FeCGT system was effective as a biocatalyst of C-glucosylation, not only for 2-hydroxyflavanones to produce flavonoid C-glucosides but also for THAP-like compounds shown in Figure 1. FeCGT_a has a relatively broad substrate preference and high enzyme activity toward these substrates (Nagatomo et al. 2014), which would have resulted in this efficient conversion of these compounds. Enzymatic conversion of phloretin to nothofagin was achieved, with a high yield, using a recombinant OsCGT coupled with sucrose synthase that could reuse the expensive UDP-glucose and enzyme (Bungaruang et al. 2013). Compared with that co-enzymatic conversion process, however, Ec-FeCGT is simpler and inexpensive because it does not require the preparation of the enzymes and the addition of the expensive UDP substrate.

In this work, we have demonstrated the production of C-glucosides of flavonoids and THAP-like compounds, using *E. coli* expressing buckwheat CGT. Similar to the concept proposed by Lim (2005) and previous works using *E. coli* cells expressing *UGT* genes (Kim et al. 2012, 2013), the glucosides produced were released into the media and were easily recovered without the need for extraction from the cells. We conclude that Ec-FeCGT is a simple and useful C-glucoside production system.

Acknowledgements

We would like to thank Ms. Sachiko Yoshioka and Dr. Tetsuya Fujimoto (Shinshu University) for the NMR analysis. We are indebted to the Division of Gene Research, Research Center for Human and Environmental Sciences, Shinshu University, for providing facilities. This work was supported in part by JSPS KAKENHI Grant No. 26450120 to G.T.

References

- Bowles D, Isayenkova J, Lim EK, Poppenberger B (2005) Glycosyltransferases: Managers of small molecules. *Curr Opin Plant Biol* 8: 254–263
- Brazier-Hicks M, Edwards R (2013) Metabolic engineering of the flavone-C-glycoside pathway using polyprotein technology. *Metab Eng* 16: 11–20
- Brazier-Hicks M, Evans KM, Gershtater MC, Puschmann H, Steel PG, Edwards R (2009) The C-glycosylation of flavonoids in cereals. *J Biol Chem* 284: 17926–17934
- Bungaruang L, Gutmann A, Nidetzky B (2013) Leloir glycosyltransferases and natural product glycosylation: Biocatalytic synthesis of the C-glucoside nothofagin, a major antioxidant of redbush herbal tea. *Adv Synth Catal* 355: 2757–2763
- Byrne PF, Darrah LL, Snook ME, Wiseman BR, Widstrom NW, Moellenbeck DJ, Barry BD (1996) Maize silk-browning, maysin content, and antibiosis to the maize earworm, *Helicoverpa zea* (Boddie). *Maydica* 41: 13–18
- de Oliveira DR, Zamberlam CR, Gaiardo RB, Rêgo GM, Cerutti JM, Cavalheiro AJ, Cerutti SM (2014) Flavones from *Erythrina falcata* are modulators of fear memory. *BMC Complement Altern Med* 14: 288
- Franz G, Grün M (1983) Chemistry, occurrence and biosynthesis of C-glycosyl compounds in plants. *Planta Med* 47: 131–140
- Gachon CM, Langlois-Meurinne M, Saindrenan P (2005) Plant secondary metabolism glycosyltransferases: The emerging functional analysis. *Trends Plant Sci* 10: 542–549
- Green MR, Sambrook J (2012) *Molecular Cloning: A Laboratory Manual*, 4th Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York
- Hooper AM, Tsanuo MK, Chamberlain K, Tittcomb K, Scholes J, Hassanali A, Khan ZR, Pickett JA (2010) Isoschaftoside, a C-glycosylflavonoid from *Desmodium uncinatum* root exudate, is an allelochemical against the development of *Striga*. *Phytochemistry* 71: 904–908
- Jay M (1994) C-Glycosylflavonoids. In: Harborne JB (ed) *The Flavonoids: Advances in Research Since 1986*. Chapman & Hall, London, UK, pp 57–93
- Jones P, Vogt T (2001) Glycosyltransferases in secondary plant metabolism: Tranquilizers and stimulant controllers. *Planta* 213: 164–174
- Jung IH, Lee HE, Park SJ, Ahn YJ, Kwon G, Woo H, Lee SY, Kim JS, Jo YW, Jang DS, et al. (2014) Ameliorating effect of spinosin, a C-glycoside flavonoid, on scopolamine-induced memory impairment in mice. *Pharmacol Biochem Behav* 120: 88–94
- Kim B, Kim H, Ahn J (2012) Production of bioactive flavonol rhamnosides by expression of plant genes in *Escherichia coli*. *J Agric Food Chem* 60: 11143–11148
- Kim H, Kim B, Ahn J (2013) Regioselective synthesis of flavonoid bisglycosides using *Escherichia coli* harboring two glycosyltransferases. *Appl Microbiol Biotechnol* 97: 5275–5282

- Lim EK (2005) Plant glycosyltransferases: Their potential as novel biocatalysts. *Chemistry* 11: 5486–5494
- Lim EK, Ashford DA, Hou B, Jackson RG, Bowles DJ (2004) *Arabidopsis* glycosyltransferases as biocatalysts in fermentation for regioselective synthesis of diverse quercetin glucosides. *Biotechnol Bioeng* 87: 623–631
- Matsuba Y, Sasaki N, Tera M, Okamura M, Abe Y, Okamoto E, Nakamura H, Funabashi H, Takatsu M, Saito M, et al. (2010) A novel glucosylation reaction on anthocyanins catalyzed by acyl-glucose-dependent glycosyltransferase in the petals of carnation and delphinium. *Plant Cell* 22: 3374–3389
- McNally DJ, Wurms KV, Labbé C, Quideau S, Bélanger RR (2003) Complex C-glycosyl flavonoid phytoalexins from *Cucumis sativus*. *J Nat Prod* 66: 1280–1283
- Nagatomo Y, Usui S, Ito T, Kato A, Shimosaka M, Taguchi G (2014) Purification, molecular cloning, and functional characterization of flavonoid C-glycosyltransferases from *Fagopyrum esculentum* M. (buckwheat) cotyledon. *Plant J*, 80: 437–448
- Ogawa K, Kawasaki A, Omura M, Yoshida T, Ikoma Y, Yano M (2001) 3',5'-Di-C- β -glucopyranosylphloretin, a flavonoid characteristic of the genus *Fortunella*. *Phytochemistry* 57: 737–742
- Talhi O, Silva A (2012) Advances in C-glycosylflavonoid research. *Curr Org Chem* 16: 859–896
- Wang LE, Bai YJ, Shi XR, Cui XY, Cui SY, Zhang F, Zhang QY, Zhao YY, Zhang YH (2008) Spinosin, a C-glycoside flavonoid from semen *Ziziphi Spinozae*, potentiated pentobarbital-induced sleep via the serotonergic system. *Pharmacol Biochem Behav* 90: 399–403
- Yepremyan A, Salehani B, Minehan TG (2010) Concise total syntheses of aspalathin and nothofagin. *Org Lett* 12: 1580–1583